

The clinical and economic value of
Mako SmartRobotics™

Mako clinical evidence



Table of contents

Mako SmartRobotics™ – an introduction.....4

Changing orthopaedic landscape
and the future of healthcare reform.....6

Overview of osteoarthritis.....6

Burden of disease6

Approaches to treatment.....6

Enhancing hip and knee arthroplasty7

The potential benefits of Mako SmartRobotics™
in total knee arthroplasty7

The potential benefits of Mako SmartRobotics™
in partial knee arthroplasty.....12

The potential benefits of Mako SmartRobotics™
in total hip arthroplasty.....17

How Mako SmartRobotics™ differs from other
robotic platforms.....21

Learning curve of Mako SmartRobotics™22

The adoption of Mako SmartRobotics™24

References.....25

List of figures

Figure 1. The Mako System..... 4

Figure 2. Challenges associated with hip and knee arthroplasty procedures that may contribute to failure or need for revision surgery..... 7

Figure 3. Study outcomes for patients who underwent conventional jig-based TKA and robotic-arm assisted TKA².....8

Figure 4. Results showed significant reduction in LOS and less MEU required for the RATKA group⁴⁰..... 9

Figure 5. FJS at two-year follow-up for RATKA patients⁴³ 10

Figure 6. Medicare 100% Standard Analytical Files were queried for RATKA and MTKA cases. Based on propensity-matched cohorts, RATKA had (a) reduced episode-of-care cost at 30,60, and 90 days postoperative as well as (b) reduced rate of readmission at those time points. Mont et al. also observed that (c) RATKA patients were more likely to be sent home postoperatively with a health aide or self-care than sent to a skilled nursing facility or inpatient rehab⁴⁹..... 11

Figure 7. Average post-index 90-day pay amounts for patients who underwent RATKA vs. MTKA⁵¹..... 12

Figure 8. Visual analog scale collected for Mako Partial Knee and manual surgery at 90 days postoperative. Data showed 55.4% lower postoperative pain for Mako Partial Knee patients compared to manual⁵⁶..... 13

Figure 9. An assessment of early functional outcomes in conventional versus robotic-arm assisted PKA⁵⁸ 13

Figure 10. Midterm patient satisfaction with medial Mako Partial Knee procedures^{3,59} 14

Figure 11. Survivorship data from Pearle et al.⁵⁹ and Kleeblad et al.³ on robotic-arm assisted PKA compared to studies in literature and annual registries reporting 2 to 3 years and 5 to 6 years conventional PKA survivorship data 14

Figure 12. Compiled revision rates demonstrating enhanced results for Mako Partial Knee^{3,19,59,61-67} 14

Figure 13. Incremental cost-effectiveness ration (ICER) values comparing PKA with NST, TKA with NST, and PKA with TKA by age⁷⁰ 15

Figure 14. Cost per quality-adjusted-life-year (QALY) of robot-assisted unicompartmental knee arthroplasty according to case volume and length of hospital stay relative to total knee arthroplasty⁷¹ 16

Figure 15. Mako vs. manual cup placement in total hip arthroplasty⁷⁸..... 17

Figure 16a and 16b. Acetabular cup placement within the Lewinnek and Callanan safe zones for Mako Robotic-Arm Assisted Total Hip Arthroplasty (A) and conventional total hip arthroplasty (B)⁸⁰..... 18

Figure 17. Postoperative outcomes for patients receiving MTHA vs. RATHA^{1,81} 20

Figure 18. Minimum five-year outcomes of robotic-assisted primary total hip arthroplasty compared to manual primary total hip arthroplasty⁸³ 20

Figure 19. Cutting robotic systems used in orthopaedics^{92,93} 22

Figure 20. Mako SmartRobotics™ statistics⁹⁹ 24

About this document

This document is intended to provide useful information to payers, healthcare facilities and healthcare providers to assess the clinical and economic value of Mako SmartRobotics™. The studies explored in this document are of varying design, ranging from large controlled clinical studies to single-surgeon studies and cadaver studies. As a result of variations in study design, the robustness of the data arising from different studies may vary. The document includes descriptions of studies relied upon, and published sources are cited throughout. We encourage you to consult the cited publications.

Mako SmartRobotics™ combines three key components, 3D CT-based planning, AccuStop™ haptic technology and insightful data analytics,* into one platform which has shown better outcomes for total hip, total knee and partial knee patients.^{1,2,3}




Mako SmartRobotics™ - an introduction

Mako SmartRobotics™ offers a transformational shift in orthopaedic practice, and ultimately in patient care, through its potential to deliver value to patients, payers and surgeons. Mako can help surgeons address the challenges of today's changing orthopaedic landscape and healthcare environment.






Figure 1. The Mako System

Know more.



It all starts with a CT scan that creates a 3D image of the patient's unique anatomy. This information allows surgeons to create their patient's plan and assess and balance the joint.

Cut less.



Using everything the CT scan allows the surgeon to know about their patient, Mako's AccuStop™ haptic technology guides the surgeon to cut what they've planned...precisely for each patient.^{1,4,5} For some patients, that means preserving soft tissue; for others, that means saving healthy bone.⁶⁻¹⁰

Throughout the procedure, surgeons and their surgical staff receive real-time data, allowing them to continually assess ligament tension throughout range of motion and implant articulation and helping them to avoid inadvertent transection of vital structures. Surgeons can refine the surgical plan intraoperatively for enhanced soft tissue balance.

Changing orthopaedic landscape and the future of healthcare reform

Demand for knee and hip procedures is on the rise. According to a study evaluating historical procedure rates and population projections using the National Inpatient Sample, primary total hip arthroplasty (THA) in the U.S. is projected to increase 71%, to 635,000 procedures, by 2030 and primary total knee arthroplasty (TKA) in the U.S. is projected to increase 85%, to 1.26 million procedures, by 2030.¹¹ These dramatic increases will have a considerable impact on healthcare utilization, demand for orthopaedic surgeons and the desire for technological advancements to enhance patient outcomes.

Overview of osteoarthritis

An estimated 22.7% (54.4 million) of adults (aged 18 years and older) have been diagnosed with arthritis in the U.S.¹² About 43.5% (23.7 million) of these 54.4 million adults have limitations in their usual activities due to their arthritis. Osteoarthritis (OA), the most common form of arthritis, is a major cause of pain and disability among adults in the U.S.¹² From 2008 to 2014, 32.5 million U.S. adults, or one in seven persons (14%), reported osteoarthritis and allied disorders, including joint pain with other specified or unspecified arthropathy, annually.¹³ Among adults 65 years and older in the U.S., an estimated 43% are living with osteoarthritis.¹³

As the U.S. population ages, the number of adults affected by osteoarthritis is expected to increase substantially.¹⁴ By the year 2040, an estimated 78.4 million (25.9% of the projected total adult population) adults will have doctor-diagnosed arthritis, and an estimated 34.6 million adults (43.2% of adults with arthritis or 11.4% of all U.S. adults) will report arthritis-attributable activity limitations.¹²

Burden of disease

The Global Burden of Disease study ranked hip and knee osteoarthritis as the 11th highest contributor to global disability.¹⁵ In the U.S., 1 in 3 adults with arthritis reports arthritis-attributable activity limitations, and the prevalence of age-adjusted arthritis-attributable social participation restriction ranges from about 1 in 8 to more than 1 in 4 adults with arthritis across states nationwide.¹² Arthritis-attributable severe joint pain is reported by at least 1 in 5 adults with arthritis in every state in the U.S.¹²

OA was the second most costly health condition treated at U.S. hospitals in 2013.¹² In that year, it accounted for \$16.5 billion, or 4.3%, of the combined costs for all hospitalizations. OA was also the most expensive condition for which privately insured patients were hospitalized, accounting for over \$6.2 billion in hospital costs.¹²

Nearly 3 million hospital stays in 2013 in the U.S. had an OA diagnosis, and it was the leading cause (46%) of hospitalization among all arthritis diagnoses. Osteoarthritis accounted for 45% of total hospital charges for arthritis diagnoses (cost charged but not necessarily paid), presumably in part because OA is the principal diagnosis associated with hip and knee joint replacements.¹³ Fewer than half (43%) of patients with an OA diagnosis were discharged to home or self-care, the lowest share of all arthritis-diagnosed hospitalized patients. This is probably due to discharges to assisted living facilities or skilled nursing facilities for rehabilitation following hip or knee joint replacement.¹³

Osteoarthritis was diagnosed in 20.8 million outpatient visits in 2013 and accounted for 1 in 5 (21%) ambulatory care visits with any arthritis diagnosis. During that time, 1 in 12 (8.4%) outpatient visits included an OA diagnosis.¹³

Combining direct and indirect costs, average annual all-cause costs for OA in the U.S. and allied disorders for the years 2008 to 2014 were \$486.4 billion. Total incremental costs (direct and indirect costs associated with osteoarthritis) were \$136.8 billion.¹³

Approaches to treatment

Joint replacement surgery is a treatment consideration for patients who are non-responsive to initial therapy and who continue to experience continuing joint symptoms and pain.¹⁶

For patients who are candidates for joint arthroplasty procedures, several surgical approaches are available, including total joint replacement and partial joint arthroplasty, as well as a variety of surgical techniques including manual (traditional), navigation-assisted and robotic-assisted techniques. While total joint replacement procedures may offer pain reduction and function recovery for many, the potential for complications still exists.¹⁷

Knee and hip arthroplasty are associated with a recovery period that may include postoperative pain, frequent physical therapy, the use of assistive devices for ambulation in the near-term and narcotic analgesics to manage pain in the months following the procedure.¹⁷

Procedure	Common challenges
Partial knee arthroplasty ¹⁸⁻²³	<ul style="list-style-type: none"> • Demanding procedure, with restricted visual field • Potential for technical errors • Poorly implanted PKA may fail earlier
Total knee arthroplasty ²⁵	<ul style="list-style-type: none"> • Instability • Infection • Aseptic loosening • Malalignment
Hip arthroplasty ²⁶	<ul style="list-style-type: none"> • Early mechanical failures • Dislocation • Prosthetic failures (periprosthetic fracture, leg length discrepancy)

Figure 2. Challenges associated with hip and knee arthroplasty procedures that may contribute to failure or need for revision surgery

In some cases, patients may be hesitant to undergo these procedures. Although many factors have been shown to influence the prevalence of knee and hip arthroplasties, patient preferences play a role in these surgeries as well. A qualitative focus-group study of ethnically and age-diverse patients with knee osteoarthritis explored factors that patients considered to be important in their decision to undergo TKA. Among these patients, personal experience (positive and negative), fear of lengthy recovery and complications, and interactions with physicians were all important decision-making factors.²⁴

Enhancing hip and knee arthroplasty

The comprehensive research on Mako has demonstrated the potential clinical, functional and economic value of the Mako System and the corresponding partial knee, total knee and total hip implant systems, and has laid a scientific foundation for the support and development of future products and applications. Studies have shown enhanced patient outcomes, reductions in health resources utilization and episode-of-care (EOC) cost-savings in PKA, TKA and THA.

The potential benefits of Mako SmartRobotics™ in total knee arthroplasty

Total knee arthroplasty is an established and successful procedure for the treatment of end-stage knee arthritis.²⁷ Survivorship at 10 years is commonly reported in the 90th percentile,²⁸ while outcomes reported using patient-reported outcome measures (PROMs) demonstrate that TKA also delivers a functional benefit to patients.²⁹

Despite the demonstrable benefits of TKA, satisfaction rates are known to be lower than for total hip arthroplasty.³⁰ Reported dissatisfaction rates for TKA are around 20%.^{31,32} TKA is also known to be sensitive to surgical factors such as implant positioning and soft tissue balance.^{33,34} Inaccuracies in positioning and soft tissue balance have the potential to reduce implant survivorship and impact negatively on patient outcomes.³³⁻³⁵

The Mako Total Knee application, in comparison to manual techniques, has been shown in cadaveric and clinical settings to have increased accuracy and precision of component placement to plan.³⁶⁻³⁸ Features of Mako SmartRobotics™ that may have contributed to these outcomes include preoperative 3D planning, which takes into account each patient’s specific anatomy, and AccuStop™ haptic technology, which enables the surgeon to execute their plan. This plan can be virtually modified intraoperatively to address implant alignment, soft tissue balancing and flexion contractures. Additional features include intraoperative visual, auditory and tactile feedback provided to the user.

Mako Total Knee was launched in 2016. As the initial Mako Total Knee patients have reached various postoperative time points, improvements in both short- and midterm outcomes have been shown. In a prospective, consecutive series, single-surgeon study, Kayani et al. demonstrated statistically significant early postoperative results for 40 patients who received Mako Total Knee surgery as compared to 40 patients who received conventional jig-based TKA.² The Mako Total Knee group had less postoperative pain ($p < 0.001$),

less need for analgesics ($p < 0.001$), less postoperative blood loss ($p < 0.001$), less time to achieve straight leg raise ($p < 0.001$), less time to hospital discharge (Mako Total Knee resulted in 26% reduction in length of stay (LOS)) and improved maximum flexion at discharge.² In summary, this study demonstrated that Mako Total Knee was associated with decreased pain, improved early functional recovery and reduced time to hospital discharge compared with conventional jig-based TKA.²

Outcome	Conventional (n=40)	Robotic (n=40)	P-value
Mean operating time (mins)	61.2 (54.6 to 83.1)	70.4 (59.2 to 91.7)	0.34*
Mean fall in Hb (g/L)	26.1 (5.1 to 49.6)	18.4 (8.0 to 37.2)	<0.001*
Mean postoperative Hb (g/L)	106.7 (77.3 to 138.4)	114.7 (86.4 to 139.1)	0.01*
Mean pain score (NRS) – Day 0	5.4 (3.0 to 7.0)	3.1 (2.0 to 5.0)	<0.001*
Mean pain score (NRS) – Day 1	6.3 (4.0 to 8.0)	3.6 (2.0 to 6.0)	<0.001*
Mean pain score (NRS) – Day 2	6.1 (3.0 to 8.0)	3.3 (1.0 to 5.0)	<0.001*
Mean pain score (NRS) – Day 3	4.5 (2.0 to 7.0)	2.6 (1.0 to 5.0)	<0.001*
Median analgesia (mg) – Day 0	36.0 (IQR 29 to 51.3)	20.0 (IQR 16.0 to 28.5)	<0.001†
Median analgesia (mg) – Day 1	10.0 (IQR 10.0 to 20.0)	10.0 (IQR 0.0 to 10.0)	<0.001†
Median analgesia (mg) – Day 2	10.0 (IQR 10.0 to 20.0)	10.0 (IQR 0.0 to 10.0)	<0.001†
Median analgesia (mg) – Day 3	10.0 (IQR 0.0 to 10.0)	0.0 (IQR 0.0 to 5.0)	<0.001†

Figure 3. Study outcomes for patients who underwent conventional jig-based TKA and robotic-arm assisted TKA²

*Unpaired t-test

†Mann-Whitney U test

NRS, numerical rating scale; IQR, interquartile range

Bhimani and colleagues published a comparison of 140 robotic-arm assisted TKA (RATKA) patients and 127 manual TKA (MTKA) patients. Consistent with Kayani et al., Bhimani et al. observed reductions in early postoperative pain, opioid usage and length of stay for patients who underwent RATKA. Patients undergoing RATKA had statistically significantly lower average visual analog scores (VAS) for pain, both at rest and with activity, at two and six weeks following the index procedure. At the six-week interval, the RATKA group required 3.2 mg less morphine equivalents per day relative to the conventional group ($p < 0.001$), and a significantly greater number of patients in the RATKA group were free of opioid use compared to the conventional TKA group (70.7% vs. 57.0% ($p = 0.02$)). Patients in the RATKA group had a shorter LOS (1.9 days vs. 2.3 days ($p < 0.001$)), and a greater percentage of RATKA patients were discharged on postoperative day one (41.3% vs. 20.5% ($p < 0.001$)).³⁹

Clark et al. published a study that compared clinical outcomes in patients who received either a haptically guided RATKA or a computer-navigated TKA (CN TKA). Compared to those who received CN TKA, patients who received RATKA had significantly improved postoperative pain, reduced total morphine consumption and a reduced length of stay. The mean LOS was 3.05 days for the RATKA group compared to 4.1 days for the CN TKA group ($p < 0.001$). There was no significant difference found between the groups in Oxford Score, Forgotten Joint Score (FJS) or EQ5D VAS at 10 weeks or one year. The authors reported a statistically significant difference in inpatient total morphine equivalent consumption, with the RATKA group at 173 units and CN TKA group at 262 units ($p = 0.001$). In addition, a positive relationship was found between morphine equivalence usage (MEU) and increase in length of stay.⁴⁰

Longer-term studies also report reduced pain and improvements in outcome scores for RATKA patients. Marchand et al. published a single-surgeon study that was performed on consecutive cemented robotic-arm assisted TKA patients matched with consecutive cemented manual TKA patients.^{41,42} In a cohort followed to six months postoperative, a WOMAC survey including pain, stiffness, and physical function subcategories was administered to patients. At six months, the RATKA cohort had significantly reduced total pain scores when compared to the MTKA cohort and also demonstrated significantly improved mean total satisfaction and physical function scores when compared to the manual cohort.⁴¹ In another cohort followed to one year postoperative, significant improvements in mean total




Figure 4. Results showed significant reduction in LOS and less MEU required for the RATKA group⁴⁰

satisfaction and physical function scores were seen when compared to the manual cohort at six months and at one year.⁴² These results indicate the potential of this surgical tool to improve short-term pain, physical function and total satisfaction scores. Although they involved limited cohorts, these studies showed promising outcomes for up to one year for RATKA patients when compared to the MTKA control group.^{41,42}

Marchand et al. continued follow-up of 196 patients longitudinally and collected two-year postoperative WOMAC, FJS and Patient Joint Perception (PJP) scores.⁴³ Patient-reported mean pain, physical function and total satisfaction scores statistically significantly improved as patients progressed from preoperative to two-year follow-up ($p < 0.05$, Figure 5). RATKA patients reported a median FJS of 65.8 ± 31.1 at two-year follow-up with 36% of patients having an FJS > 80 . The median FJS was comparable to the normative value, 66.8 ± 34.0 , reported for a U.S. general population with a similar age range.⁴³ Based on the PJP score, 83% of patients reported their knee feeling like a “natural joint” or an artificial joint with minimal or no restrictions.




Figure 5. FJS at two-year follow-up for RATKA patients⁴³

Wang et al.⁴⁴ performed a retrospective review in which a single high-volume surgeon performed 148 RATKA cases and 159 MTKA cases with matched demographics. The RATKA cohort experienced a significantly longer tourniquet time when the learning curve phase was included (96.8 min vs. 91.6 min); however, a significant difference was not observed when the last 20 RATKA cases were compared to the MTKA cases (93.8 min vs. 91.6 min, $p = 0.506$). Postoperatively, the RATKA cohort was more often discharged to home care (95.95% vs. 83.65%, $p < 0.001$) compared to acute rehabilitation, had a reduced number of physical therapy appointments (11.0 vs. 13.3, $p = 0.004$) and a lower number of 30-day readmissions (1 vs. 5, $p = 0.014$). This trend in enhanced outcomes followed through to one year, with the RATKA group demonstrating enhanced Knee Injury and Osteoarthritis Outcome Scores for Joint Replacement (KOOS-JR) ($p = 0.034$) and FJS ($p = 0.021$). These favorable results for the RATKA indicate patient outcomes continued to improve when compared to MTKA out to one year postoperative.

As more robotic-arm assisted TKA patients reach two-year follow-up, additional studies are beginning to report on these outcomes. Malkani et al. reported on the two-year outcomes of 188 RATKAs performed at five centers. They found that RATKA patients had excellent outcomes in multiple PROMs. The mean postoperative Short Form-12 Questionnaire (SF-12) mental composite score (MCS) and physical composite score (PCS) were both 57 points, with 50 as the threshold for norm-based scoring. The mean FJS was 75 points. The mean Knee Society Score (KSS) Functional Score was 84 points and the mean Knee Score was 92 points. Malkani et al. also found that the aseptic revision rates were low at 1.06% and that there were few other postoperative complications (3.7%).⁴⁵ A separate analysis on the manipulation under anesthesia (MUA) rates of these

patients compared with a consecutive equal number of control patients by each of the specific surgeons found that patients who underwent robotic-assisted TKA experienced a significant 4.5-fold decrease in rates of manipulation under anesthesia ($p = 0.032$). Given that MUAs can be a marker of knee stiffness following total knee arthroplasty, the lower rate indicates that study cohort patients had less knee stiffness and, therefore, greater initial postoperative range of motion than the control cohort.⁴⁶

Mako Total Knee provides surgeons with preoperative planning and real-time data, allowing for continuous assessment of ligamentous tension and range of motion. Using this technology, soft tissue protection,^{8,47} reduced early postoperative pain,² improved patient satisfaction,⁴² reduced complications such as MUAs,⁴⁶ and reduced LOS³⁹ have been shown. These advances have the potential to enhance surgical outcomes and may also reduce episode-of-care costs for patients, payers and hospitals. As Mako SmartRobotics™ continues to be adopted, it is also important to understand whether Mako Total Knee is associated with reduced EOC costs.


Cool et al. performed a retrospective review of a U.S.-based Medicare database for TKA surgeries between January 2016 and March 2017.⁴⁸ After propensity score matching, 519 RATKA and 2,595 MTKA cases were assessed to compare EOC cost, index cost, LOS, discharge disposition and readmission rates. Results found overall 90-day EOC costs were \$2,391 less for RATKA patients ($p < 0.0001$).⁴⁸ Index facility cost and LOS were less for RATKA patients by \$640 ($p = 0.0001$) and 0.7 days ($p < 0.0001$), respectively.⁴⁸ Additionally, robotic-arm assisted patients were discharged to self-care more frequently (56.65% vs. 46.67%, $p < 0.0001$) and to skilled nursing facilities (SNF) less frequently (12.52% vs. 21.70%, $p < 0.0001$), and had a 90-day readmission reduction of 33% ($p = 0.04$) compared to MTKA patients.⁴⁸ This evidence demonstrated a cost-savings to Medicare when comparing RATKA to MTKA. This 90-day EOC savings for the RATKA group was driven by reduced facility costs, LOS and readmissions, and an economically beneficial discharge destination.⁴⁸

Mont et al. performed a healthcare utilization analysis that compared RATKA and MTKA techniques.⁴⁹ They specifically compared (1) index costs and (2) discharge dispositions, as well as (3) 30-day (4) 60-day and (5) 90-day (a) episode-of-care costs, (b) readmission, and (c) postoperative healthcare utilization. The same propensity matched group from Cool et al. was used in


this study to assess trends in total episode payments, healthcare utilization and readmissions at 30-, 60- and 90-day time points. The RATKA cohort had consistently lower average total episode payment than the MTKA cohort when compared at 30, 60, and 90 days (Figure 6). At 30 days, 47% fewer RATKA patients utilized SNF services (13.5 vs. 25.4%, $p < 0.0001$, Figure 6) and RATKA patients had lower SNF costs at 30, 60, and 90 days. Robotic-arm assisted TKA patients also utilized fewer home health visits and incurred fewer costs at each time point ($p < 0.05$). Additionally, 31.3% fewer RATKA patients utilized emergency room services at 30 days postoperatively, and the RATKA cohort had fewer 90-day readmissions (5.2 vs. 7.75%, $p = 0.0423$, Figure 6). Mont et al. concluded that RATKA was associated with lower 30-, 60- and 90-day postoperative costs and healthcare utilization. These results are of marked importance given the emphasis to contain and reduce healthcare costs, and this study provides promising initial economic insights into RATKA.

While total joint arthroplasties account for more Medicare expense than any other inpatient procedure,⁵⁰ studies have reported the growth of TKA procedures in commercially insured patients under 65. Pierce and colleagues⁵¹ evaluated 90-day EOC costs in a commercially insured population. TKA procedures were identified using the Optum Insights Inc. database. The procedures were stratified into two groups, the RATKA cohort or the MTKA cohort. Following 1:5 propensity score matching, 357 RATKAs and 1785 MTKAs were included in the analysis. Utilization and associated costs were analyzed for 90 days following the index procedure. The authors observed that the overall length of stay was significantly lower for those in the RATKA arm (1.80 vs. 2.72 days; $p < 0.0001$). Within the 90 days following the index stay, patients who underwent robotic-arm assisted TKA were less likely to utilize inpatient services (2.24 vs. 4.37%; $p = 0.0444$) or SNF (1.68 vs. 6.05%; $p < 0.0001$) than those in the MTKA cohort. Patients who utilized home health in the RATKA arm used significantly fewer days of home health than MTKA patients (5.33 vs. 6.36 days; $p = 0.0037$). Cost associated with the utilization of services was substantially lower in the RATKA arm; the overall post-index cost was \$1,332 less per case in the RATKA arm (\$6,857 vs. \$8,189; $p = 0.0018$). Cost was also significantly less in the RATKA cohort for those patients who utilized outpatient rehab (\$2,272 vs. \$2,494; $p = 0.0194$) and pharmacy (\$588 vs. \$843; $p = 0.0057$). The 90-day EOC cost was \$4,049 less per case in the RATKA arm (\$28,204 vs. \$32,253; $p < 0.0001$).⁵¹

a: Total episode costs at 30, 60 and 90 days: RATKA vs. MTKA



b: Readmission rates at 30, 60 and 90 days: RATKA vs. MTKA



c: Discharge destination




Figure 6. Medicare 100% Standard Analytical Files were queried for RATKA and MTKA cases. Based on propensity-matched cohorts, RATKA had (a) reduced episode-of-care cost at 30, 60, and 90 days postoperative as well as (b) reduced rate of readmission at those time points. Mont et al. also observed that (c) RATKA patients were more likely to be sent home postoperatively with a health aide or self-care than sent to a skilled nursing facility or inpatient rehab compared to manual⁴⁹

Average post-index 90-day pay amounts




Figure 7. Average post-index 90-day pay amounts for patients who underwent RATKA vs. MTKA⁵¹

*indicates statistically significant difference

The potential benefits of Mako SmartRobotics™ in partial knee arthroplasty

Stryker’s Mako SmartRobotics™ technology is designed to help enhance the accuracy of component placement, as well as the reproducibility of partial knee arthroplasty.

Partial knee resurfacing for patients with osteoarthritis isolated to only one or two compartments is designed to spare the anterior and posterior cruciate ligaments as well as healthy bone and tissue. Minimizing tissue disruption may enhance patient outcomes and recovery time after TKA procedures, thereby reducing the risk of complications, associated costs and hospital days.^{52,53} Manual partial knee replacement can be a demanding procedure with a restricted field of view, and surgeons cannot preoperatively create a patient-specific plan.⁴ Patellofemoral arthroplasty is a particularly challenging procedure due to the need to place components properly in multiple planes. This procedure can be sensitive to even one millimeter of abnormality in implant depth, and poorly implanted components may fail earlier.⁴ With manual instrumentation, it can be difficult to consistently restore tibial slope,¹⁸ coronal alignment, femoral rotation and limb alignment.⁵⁴

A key clinical paper on Mako accuracy, published by Bell et al., reported on a randomized controlled trial (RCT) involving 120 patients. The study compared patients who received robotic-arm assisted PKA (Restoris MCK n=62) with those who underwent manually implanted PKA (Oxford n=58).⁴ Comparisons were made between groups in terms of the preoperative plan of femoral and tibial component positioning against the actual alignment achieved in three different planes (axial, coronal and sagittal). Results showed more accurate component positioning in the robotic-arm assisted group, with lower root mean square errors and significantly lower median errors in all six component parameters (p < 0.01).⁴ The proportion of patients with tibial slope within 2° of the target position was significantly greater using the robotic-arm assisted technique than the manual technique (80% compared with 22%, p = 0.0001). It was concluded that the Mako System helped surgeons to more consistently place the PKA implant in accordance with the preoperative plan.⁴

These results were corroborated by a study performed at University College Hospital in London, England, by Kayani et al.⁵⁵ A single surgeon compared implant placement accuracy using radiographs from 60 consecutive conventional PKAs (Oxford) compared to the surgeon's first 60 consecutive Mako Partial Knees (Restoris MCK). The Mako group had significantly ($p < 0.001$) more accurate placement to plan of the femoral and tibial implants, as well as more accurate recreation of the knee's mechanical alignment, posterior tibial slope and joint line height.⁵⁵

Outcomes for partial knee arthroplasty

Achieving desired alignment during surgery may result in enhanced outcomes and patient functioning. In a prospective, randomized, controlled single-center blinded trial ($n=139$ patients), patients were randomized to receive either a manual PKA or a Mako Partial Knee. An analysis of the RCT patients found that patients who underwent medial Mako Partial Knee experienced less pain during the 90-day postoperative period than those who underwent manual surgery.⁵⁶ Median pain scores were 55.4% lower in robotic-arm assisted patients compared to manual patients from day one to day 56 (Figure 8).⁵⁶ Furthermore, the robotic-arm assisted patients had better American Knee Society Scores (AKSS) at three months postoperatively and one year postoperatively, and a greater proportion of robotic-arm assisted patients showed improvements in their UCLA Activity Scores.⁵⁶




Figure 8. Visual analog scale collected for Mako Partial Knee and manual surgery at 90 days postoperative. Data showed 55.4% lower postoperative pain for Mako Partial Knee patients compared to manual⁵⁶

Additionally, the proportion of patients who achieved an FJS of $> 80\%$ at three months postoperatively was almost double in the robotic-arm assisted cohort compared to the manual PKA cohort, although there was no overall statistical difference.⁵⁶ The authors also found that inpatient length of stay was shorter in the robotic-arm assisted surgery group, with a difference of 0.54 days ($p = 0.07$), and observed that at three months postoperatively, primary care utilization (calculated from the proportion of the group who visited their general practitioners) was 15% lower ($p = 0.092$) in the robotic-arm group. These patients were followed out to two years postoperative and the Mako Partial Knee patients demonstrated 100% survivorship at two years postoperative, compared to 96.3% in the manual group.⁵⁷

Another study compared a consecutive series of 73 Mako Partial Knee patients with 73 manual PKA patients and found Mako Partial Knee patients to have lower postoperative pain scores at each day of hospitalization following surgery, shorter lengths of stay, reduced usage of postoperative analgesia and fewer physiotherapy (PT) visits required to achieve PT goals.⁵⁸




Figure 9. An assessment of early functional outcomes in conventional versus robotic-arm assisted PKA⁵⁸

Mako Partial Knee has also shown improvements in patient satisfaction. In a multicenter, longitudinal clinical trial, the vast majority of patients were “very satisfied” or “satisfied” with their joint replacement (Figure 10).^{3,59} This study performed follow-up at 2.5 years (909 knees) and 5.5 years (432 knees) with patients who underwent medial Mako Partial Knee procedures and a total of 92% of patients reported satisfaction with their knee 2.5 years postoperatively, while 91% of patients reported satisfaction at 5.5 years.^{3,59}

Mako Partial Knee patient satisfaction




Figure 10. Midterm patient satisfaction with medial Mako Partial Knee procedures^{3,59}

In a separate study, Zuiderbaan et al. administered the Forgotten Joint Score questionnaire to medial Mako Partial Knee patients and manually instrumented TKA patients one and two years postoperatively. Scores were compared between 65 patients who underwent medial Mako Partial Knee and 65 patients who underwent manually instrumented TKA.⁶⁰ Results demonstrated patients who underwent medial robotic-arm assisted PKA were more likely to forget their artificial joint in daily life.⁶⁰

Survivorship in partial knee arthroplasty

A multitude of studies have shown low revision rates for Mako Partial Knee. A multicenter, longitudinal study evaluating short and midterm survivorship of robotic-arm assisted medial PKA demonstrated 98.8% survivorship (in 909 knees) at 2.5-year follow-up and 97% (in 432 knees) at 5.5-year follow-up.^{3,59} More recently, five-year follow-up of 845 patients (1018 knees) has shown survivorship for medial onlay at 97.8%, lateral unicompartmental knee arthroplasty (UKA) at 97.7% and patellofemoral arthroplasty / bicompartamental knee arthroplasty at 93.3%.¹⁰² These survivorship rates were greater than rates derived from high-volume surgeon data and registry data for conventional PKA.^{3,59} The study concluded that the favorable survivorship observed resulted from Mako’s ability to help enable surgeons to achieve more accurate component positioning when compared to implant placement using manual techniques.^{3,59}

Partial knee survivorship




Figure 11. Survivorship data from Pearle et al.⁵⁹ and Kleebblad et al.³ on robotic-arm assisted PKA compared to studies in literature and annual registries reporting 2 to 3 years and 5 to 6 years conventional PKA survivorship data




Figure 12. Compiled revision rates demonstrating enhanced results for Mako Partial Knee^{3,19,59,61-67}

Similar promising data was published in the 2019 Australian Joint Registry, which reported the cumulative revision rate for the Restoris MCK medial PKA as 1.5% at one year and 2.5% at 3 years, which was significantly lower than non-robotic UKAs in the registry. This compared favorably to the revision rate for all Oxford medial PKA replacements at one year (2.2%) and at three years (5.8%) and is the lowest rate for any PKA implant reported.^{19,68} The revision rates for Mako Partial Knee have been published in cohort studies, economic analyses, level I clinical trials (RCTs) and international registries. The evidence supports excellent survivorship of the Restoris MCK implant when used with the Mako System. In summary, Mako has demonstrated positive outcomes through more accurate component positioning⁴ and high patient satisfaction.^{3,59}

Mako Partial Knee health economics

Clinical findings such as reduced revision rates have the potential to add value in the continuum of care. In a study by Cool et al., reasons for revisions and associated costs were analyzed for robotic-arm assisted partial knee arthroplasty cases. PKA procedures were identified using a U.S. commercial administrative claims database to evaluate hospital admissions for revision surgeries. Robotic-arm assisted PKA (RAPKA) and manual PKA (MPKA) procedures performed between March 1, 2013 and July 31, 2015 were used to calculate the rate of revisions within 24 months of the index procedure. Cases were propensity matched 2:1 based on age, sex, race, geographic division, high-cost comorbidities and concentration of healthcare specialists per 100,000 population to control for outside confounding factors at case index. A total of 738 commercial health plan patients (246 RAPKAs, 492 MPKAs) were selected for inclusion in the analysis. Results indicated fewer revision procedures in RAPKA, 0.81% [2/246] vs. 5.28% [26/492]; ($p = 0.0017$) and RAPKA patients incurred lower mean costs for the index stay plus revision(s), \$26,001 vs. \$27,977; $p > 0.05$. Lower length of stay at index was also noted in the RAPKA group, 1.77 vs. 2.02 days; $p = 0.0047$. The study concluded that patients who underwent RAPKA had fewer revision procedures, shorter LOS and incurred lower mean costs at 24 months.⁶⁴

Some have tried to evaluate potential clinical and economic differences between PKA and TKA. A prospective study of 30 Mako Partial Knees compared to 90 propensity-matched manual TKAs found that six-month pain VAS scores, Oxford Knee Score (OKS), FJS

and EQ5D were significantly better for the Mako Partial Knee group compared to manual TKA. They also found that LOS was significantly shorter in the robotic-arm assisted PKA group compared to manual TKA.⁶⁹

With rising demand for PKA in patients who seek restored function and a quicker recovery time, a study performed by Kazarian et al. evaluated the cost-effectiveness of PKA compared to TKA as well as nonsurgical treatment (NST). Using a Markov decision analytic model, the authors assessed lifetime costs and quality of life years (QALYs) as a function of age at time of initial treatment (ATIT) of patients with end-stage unicompartmental knee osteoarthritis. The analysis included direct medical and indirect costs. Models were run for ATITs at five-year intervals from 40 through 90 years of age. Results indicated unicompartmental knee arthroplasty had the greatest QALY accumulation followed by TKA and NST, and PKA was more cost-effective compared to NST for patients from ages 40 to 86. When surgical treatments were compared, PKA dominated TKA by generating more QALYs than TKA for all ATITs. The authors further concluded that if PKAs were performed as 12% to 20% of the total volume of knee arthroplasties versus the less than 8% observed, it would lead to a lifetime cost-savings of 987 million to 1.5 billion U.S. dollars and increased lifetime QALY accumulation of 124,403 to 217,705 across the U.S. population.⁷⁰




Figure 13. Incremental cost-effectiveness ratio (ICER) values comparing PKA with NST, TKA with NST, and PKA with TKA by age⁷⁰

In a separate U.K.-based study, a Markov decision analysis was performed to assess the cost-effectiveness of robotic-arm assisted unicompartmental knee arthroplasty (RAUKA) relative to manual TKA and PKA for patients with isolated medial compartment OA of the knee with a mean age of 65 years. The study objective was to identify the cost per QALY of RAUKA specifically relative to TKA and PKA. Model inputs included hospital costs, implant survival and mortality rate. Using a model with an annual case volume of 100 patients, the cost per QALY of RAUKA was £1395 and £1170 relative to TKA and PKA, respectively. The cost per QALY was influenced by case volume: a low-volume center performing ten cases per year would achieve a cost per QALY of £7,170 and £8,604 relative to TKA and PKA. For a high-volume center performing 200 RAUKAs per year with a mean two-day length of stay, the cost per QALY would be £648; if performed as day cases, the cost would be reduced to £364 relative to TKA. For a high-volume center performing 200 RAUKAs per year with a shorter length of stay of one day relative to PKA, the cost per QALY would be £574. Furthermore, the cost per QALY of RAUKA decreased with reducing length of hospital stay and with increasing case volume compared with TKA and PKA.⁷¹ The model showed that RAUKA may offer a cost-effective alternative to TKA and PKA for patients




Figure 14. Cost per quality-adjusted-life-year (QALY) of robot-assisted unicompartmental knee arthroplasty according to case volume and length of hospital stay relative to total knee arthroplasty⁷¹

	RAUKA ⁷¹	TKA ⁷¹	MUKA ⁷¹
Total health gain (QALYs) ⁷¹	13.59	11.8	12.2
Health improvement (vs. TKA) ⁷¹	1.8	0	1.39
Cost/QALY (vs. TKA) ⁷¹	£1,395.00	£2,101.00	£1,170.00

with isolated medial compartment OA of the knee. In summary, these models demonstrated that PKA can be more cost-effective than nonsurgical treatment and TKA for the specified age groups modeled and showed that robotic-arm assisted PKA can be cost-effective compared to TKA.

Robotic-arm assisted PKA procedures may also provide value for hospitals. A hospital in Brisbane, Australia examined the potential cost-savings for the health system and the community through the increased utilization of PKA using robotic-arm assisted PKA vs. conventional TKA. They retrospectively reviewed 240 patients where the first 120 consecutive Mako Partial Knees performed during this period were matched to 120 conventional TKAs. Clinical data from the medical records and costs for procedure for each component were collected. Bivariate analyses were performed on the data to determine if there were statistically significant differences by surgery type in clinical outcomes and financial costs. The hospital found a

significantly lower cost incurred for RAPKA vs. TKA with an average saving of AU\$7,179 per case. The operating time (86.0 min vs. 75.9 min; $p = 0.004$) was significantly higher for RAPKA compared to TKA but the length of stay was significantly lower (1.8 vs. 4.8 days; $p < 0.001$). This study also found a significant difference in the use of opioids with RAPKA compared to TKA (125.0 morphine equivalent (ME) vs. 522.1 ME, $p < 0.001$).⁷²

Studies comparing PKA to TKA have observed that PKA typically requires less rehabilitation,⁷³ results in fewer postoperative complications,⁷⁴ results in patients more likely to forget their artificial joint in daily life⁶⁰ and results in improved quality of life.⁷⁰ Studies of Mako Partial Knee have not only demonstrated improvements in short-term outcomes^{56,58} compared to manual PKA, but have also shown more favorable revision rates^{3,19,59,61-67} compared to manual PKA and demonstrated revision rates similar to those seen in TKA. These observed clinical outcomes, coupled with the potential cost-savings demonstrated in assessments of cost-effectiveness, show that Mako Partial Knee has the potential to offer both short- and long-term advantages to patients, providers and payers.

The potential benefits of Mako SmartRobotics™ in total hip arthroplasty

Component positioning (stability and dislocation)

Total hip arthroplasty has been one of the most successful procedures within the field of orthopaedics since the late 1960s.⁷⁵ The short and long-term outcomes of THA may be influenced by several factors, including patient demographics, surgical technique and implant features.⁷⁶ One of the most important surgeon-controlled factors is component positioning.⁷⁶ Component malposition has been linked to higher rates of hip dislocations, poor biomechanics, accelerated wear, leg length discrepancy and revision surgeries.⁷⁶ In addition, component malposition is directly associated with dislocations and mechanical loosening, which account for approximately 40% of THA revisions.⁷⁷ Successful clinical outcomes following total joint replacement are dependent on component placement and on restoring the natural joint anatomy of the hip.⁷⁶ Instability, early mechanical failures and dislocation in hip arthroplasty continue to be primary reasons for revision.⁷⁶

The Mako System is designed to help the surgeon minimize the margin of error associated with component placement and to enhance the accuracy and reproducibility of THA. In a U.S. multicenter clinical trial including 110 patients, acetabular cup position was compared between preoperative plan, intraoperative assessment and achieved radiographic measure. Results confirmed that surgeons using intraoperative robotic-arm assistance achieved greater accuracy to plan in preparation and position of the acetabular cup during THA.⁷⁸ Consecutive primary robotic-arm assisted THAs (RATHAs) performed by one surgeon at three intervals were analyzed in a retrospective cohort study: the initial 100 consecutive manual THAs (MTHAs) in clinical practice (2000), the last consecutive 100 MTHAs before RATHA technology introduction (2011), and the first consecutive 100 RATHA cases (2012). The rate of acetabular component placement within the Lewinnek safe zone was the highest in the RATHA cohort (77%), followed by late MTHA (45%) and early MTHA (30%). RATHA resulted in an additional 71% improvement in accuracy to plan in the first year of use.¹




Figure 15. Mako vs. manual cup placement in total hip arthroplasty⁷⁸

In another study involving six surgeons at a single U.S. institute, 1,980 THA surgeries were evaluated. The aim of this study was to understand the influence of surgical approaches and modes of guidance. Robotic-arm assisted THA resulted in a significantly greater percentage of components placed in Callanan safe zones than all other modalities, including navigation- and fluoroscopy-guided approaches.⁷⁹

In a study conducted between 2008 and 2012 comparing THA using manual alignment techniques with THA using Mako robotic-arm assisted alignment, Mako Total Hips were matched to historical manual THAs.⁸⁰ As shown in Figure 16a, 100% of the robotic-arm assisted

THAs were placed within the Lewinnek safe zone for anteversion and inclination vs. 80% (40/50) of the manually aligned and implanted THAs in Figure 16b. Similarly, 92% of the RATHAs were within the Callanan safe zone vs. only 62% of MTHAs.⁸⁰




Figure 16a and 16b. Acetabular cup placement within the Lewinnek and Callanan safe zones for Mako Robotic-Arm Assisted Total Hip Arthroplasty (A) and conventional total hip arthroplasty (B)⁸⁰

Mako Total Hip clinical success

Mako Total Hip has demonstrated:

- Enhanced acetabular component placement **accuracy** and **reduced dislocation rates** and **blood loss** when compared with MTHA¹
- Favorable short-term patient-reported outcomes^{81,82,83}
- The **highest Forgotten Joint Score** reported in literature for THA⁸²
- Decreased length of stay compared to MTHA⁸⁴

Potential to restore leg length and hip biomechanics (offset)

Manual total hip procedures may be associated with discrepancies in leg length following surgery.⁸⁵ A study examined two methods of intraoperative leg length assessment and found that a discrepancy in leg length of fewer than five millimeters was achieved in only 73% and 67% of patients for the two methods individually. The same study observed that 25% of patients had a leg length discrepancy of more than five millimeters regardless of which manual surgical method was used.⁸⁵

Another study by Manzotti et al. found that at six months postoperative, the mean postoperative leg length discrepancy was reduced to 5.06 mm (range: 0–12) in a computer-assisted group, compared to 7.64 mm (range: 0–20) in the freehand group.⁸⁶ Harris Hip Scores (HHS) post-THA have been reported to be significantly higher in patient groups in which femoral offset was normal or increased relative to the contralateral side.⁸⁷

The use of Mako Total Hip has demonstrated accuracy in achieving desired leg length. In a prospective study, 20 patients received Mako Total Hip and had postoperative CTs performed to measure accuracy to plan of acetabular and femoral implant placement. Postoperative measurements reported accurate recreation of the overall hip length and offset (1.6 mm, standard deviation (sd) 2.9 mm and 0.5 mm, sd 3.0 mm, respectively). Mean stem version as well as mean shell anteversion and inclination angles were similar between intraoperative and postoperative measurements.⁸⁸ In a two-year follow-up study of 162 Mako Total Hip patients performed by a single surgeon, no leg length discrepancies were observed.⁸²

Allowance for preservation of acetabular bone stock

Preservation of acetabular bone during primary THA is important since proper implant stability and longevity depend largely on the amount of bone stock left after acetabular reaming.⁷ Eccentric or excessive acetabular reaming may lead to soft tissue impingement, loosening, altered center of rotation, bone-to-bone impingement, intraoperative periprosthetic fracture, early implant failure due to lack of bone ingrowth and other complications, potentially leading to subsequent revision of THA.⁷ In a matched-pair controlled study, the size of the acetabular cup relative to that of the femoral head was used as a surrogate measure of acetabular bone resection. In this study, Mako Total Hip allowed for the use of smaller acetabular cups in relation to the patient's femoral head size compared to conventional THA, indicating greater preservation of bone stock.⁷

Outcomes for total hip arthroplasty

Based on data prospectively collected on primary THAs conducted since August 2000 from a single institution, Mako Total Hip was associated with enhanced accuracy and reproducibility of component placement and reduced early dislocation rates compared to conventional THA as discussed above.^{1,81} In this analysis, data was reviewed for all THAs (n = 300 patients) conducted by one fellowship-trained surgeon at a single institution over three time periods in order to compare surgical outcomes^{1,81}:

- Group one (2000 to 2001): First 100 consecutive MTHA cases conducted
- Group two (2011): Last 100 consecutive MTHA cases conducted
- Group three (2011 to 2012): First 100 consecutive Mako THA cases conducted

As shown in Figure 17, Mako Total Hip demonstrated greater accuracy for both acetabular abduction (AAB) and acetabular anteversion (AAV) and demonstrated lower dislocation rates at one year compared with manual THA.^{1,81} The average estimated blood loss was also reduced in the patient group that received robotic THA compared to manual.^{1,81}

	First 100 manual THA cases	Last 100 manual THA cases	First 100 Total Hip cases
Early dislocation rate (within first 12 months postoperative)	5%	3%	0%
Limb length discrepancy > 1.5 cm	10%	1%	1%
Estimated blood loss	534 mL	438 mL	358 mL
AAB in target zone	66%	91%	100%
AAV in target zone	39%	48%	77%
AAB and AAV in target zone	30%	45%	77%

Figure 17. Postoperative outcomes for patients receiving MTHA vs. RATHA^{1,81}

In this same study, while excellent clinical outcomes were noted for both Mako Total Hip and MTHA at a one-year clinical follow-up, patients who had received Mako Total Hip demonstrated significantly higher modified Harris Hip Scores and UCLA activity level compared with MTHA.^{1,81}

In a U.S. single-surgeon prospective study of 162 robotic-arm assisted THA patients with minimum follow-up of two years, the mean Forgotten Joint Score-12 (FJS-12), a patient-reported outcome instrument developed to assess the patient’s ability to forget the artificial joint in everyday life, was 83.1. This was the highest FJS-12 score ever reported in publicly available literature.⁸²

More recently, Domb and colleagues published five-year outcomes of 66 RATHAs propensity matched with 66 MTHAs. They found that the RATHA group reported significantly higher Harris Hip Score, Forgotten Joint Score-12, Veterans RAND-12 Physical, and 12-Item Short Form Survey Physical (P = 0.001, P = 0.002, P = 0.002, P = 0.001, respectively). The acetabular implant placement by surgeons performing RATHA had a 9- and 4.7-fold reduced risk of placement outside the Lewinnek and Callanan safe zones, respectively (relative risk, 0.11 [95% confidence interval, 0.03 to 0.46]; P = 0.002; relative risk, 0.21 [95% confidence interval, 0.01 to 0.47]; P = 0.001). In addition, RATHA recipients had lesser absolute values of leg length discrepancy and global offset (P = 0.091, P = 0.001). This study demonstrated favorable outcomes for RATHA compared to MTHA at five years postoperative.⁸³

Patient-reported outcomes	Robotic-assisted THA	Manual THA	p-value
HHS	90.57±13.46	84.62±14.45	<0.001
FJS-12	82.69±21.53	70.61±26.74	0.002
VAS	1.27±2.20	1.07±1.87	0.45
Satisfaction	8.91±2.00	8.52±2.62	0.35
VR-12 mental	60.76±5.94	58.97±6.93	0.17
VR-12 physical	50.30±8.83	45.92±9.44	0.002
SF-12 mental	56.59±5.60	56.20±6.62	0.81
SF-12 physical	48.97±9.21	44.01±10.26	0.001

Figure 18. Minimum five-year outcomes of robotic-assisted primary total hip arthroplasty compared to manual primary total hip arthroplasty⁸³

Mako Total Hip health economics

In a Medicare analysis of the 90-day episode-of-care cost of 938 RATHAs propensity matched to 4,670 MTHAs, RATHA patients were less likely to have post-index IPR or SNF admissions (0.64% vs. 2.68%; $p < 0.0001$ and 20.79% vs. 24.99%; $p = 0.0041$, respectively). RATHA patients used fewer days in post-index inpatient and SNF care (7.15 vs. 7.91; $p = 0.8029$ and 17.98 vs. 19.64; $p = 0.5080$, respectively) and used fewer HHA visits (14.06 vs. 15.00; $p = 0.0006$) compared to MTHA. RATHA had lower costs for: IPR (\$11,490 vs. \$14,674; $p = 0.0470$), SNF (\$9,184 vs. \$10,408, $p = 0.0598$) and HHA (\$3,352 vs. \$3,496; $p = 0.0133$) compared to MTHA. Overall, RATHA patients had 12% (\$948) lower average post-index costs compared to MTHA patients ($p = 0.0004$). Total 90-day EOC costs for RATHA patients were found to be \$785 less than that of MTHA patients (\$19,734 vs. \$20,519, $p = 0.0095$).⁸⁹

In summary, use of the Mako System in total hip arthroplasty has demonstrated more accurate component positioning,^{1,80,81} bone preservation,⁷ improved clinical outcomes and the potential for cost-savings.⁸⁹



How Mako SmartRobotics™ differs from other robotic platforms

Mako SmartRobotics™ possesses several key features that differentiate it from other robotic surgery platforms. After a thorough surgical plan is created and approved by the surgeon, the Mako System assists surgeons with executing that plan using AccuStop™ haptic technology. The implant position, tracking and soft tissue balancing are assessed in a virtual 3D model by combining a preoperative CT and intraoperative bone registration. A CT scan uses a combination of 2D and digital geometry processing to generate a 3D image of the body. While plain film radiographs (X-rays) provide a 2D image of the scanned area, anatomic structures may overlap, creating an image which is less detailed than a CT scan. In a CT image, overlapping structures are eliminated, making the internal anatomy easier to visualize. In knee and hip arthroplasty procedures, the femoral version and tibial torsion⁷⁶ can provide critical guidance when planning a case. Bony anatomic landmarks of the femur and tibia can be clearly identified using 3D imaging technologies. After a surgeon assesses implant size and position in the preoperative plan, the robotic arm is introduced to the surgical site. The robotic arm uses AccuStop™ haptic technology to help ensure only the desired bone is resected. The robotic arm will give resistance, an audible warning and ultimately turn off if the surgeon attempts to move the cutting tool on the robotic arm outside the boundaries created in the preoperative plan.

While various other robotic systems include a robotic arm, robotic-guided cutting jigs or different navigation strategies,⁹⁰ the Mako System has the capability to virtually create and modify the 3D preoperative plan before an incision is made, and the surgeon is able to analyze and modify the preoperative plan before bone resection even begins.

Some joint arthroplasty techniques do not require a CT scan at all prior to surgery. For example, some orthopaedic navigation systems are “model-based”, where information from a CT scan is not utilized. Instead, navigation software calculates an individual model of the patient’s anatomy based on defined landmarks on the bone, which are acquired using a navigated instrument (registration).⁹¹ After optional planning on the model (e.g., virtual orientation and placement of the joint implant), the actual procedure follows, where the surgeon is supported by relevant information added through the navigation system (navigation).⁹¹

Surgeons may not adopt robotic technology and may instead elect to continue to perform their cases manually. For manual total knee arthroplasty, a surgeon uses X-rays of the joint to visually identify the desired implants and positioning/alignment of the implants. During the surgery, mechanical instruments such as rods placed inside or outside of the bones and blocks are used to measure and assess the angle and resection depth of the bone cuts. The bone cuts are performed with a hand-held powered saw, which is typically guided by a cutting block which has been pinned to the bone. This technique requires the surgeon to be able to visualize the

edges of the bone while making the cuts in order to avoid cutting into the soft tissues inadvertently. The surgeon then uses trial implants to assess the cuts and make any alterations necessary before the final implants are placed and the wound is closed.

Mako is the only robotic system that has demonstrated, through published clinical studies, higher accuracy and precision to plan for implant placement and sizing for total knee, total hip and partial knee arthroplasty compared to manual techniques.^{4,79,94} Mako is also currently the only robotic-arm bone preparation system in the marketplace that uses AccuStop™ haptic technology and that has the ability to cut with a saw, burr with a burr and ream with a reamer.

Learning curve of Mako SmartRobotics™

The learning curves of robotic-arm assisted TKA, PKA and THA have been explored in the literature. Kayani and colleagues evaluated 60 Mako Partial Knees and compared them to 60 manual partial knees. They found their learning curve for surgical time and surgical team confidence levels to be six cases. They also found that improved accuracy to plan was experienced from the first case, indicating that Mako Partial Knee surgery did

Figure 19. Cutting robotic systems used in orthopaedics^{92,93}

System*	Application†	Cutting type	Cutting control
TSolution One	TKA	Direct	Autonomous
Mako	UKA	Direct	Haptic
Mako	TKA	Direct	Haptic
Mako	THA	Direct	Haptic
NAVIO	UKA	Direct	Boundary control
NAVIO	TKA	Indirect	Boundary control
ROSA	TKA	Indirect	Cutting guide
OMNIBotics	TKA	Indirect	Cutting guide
SpineAssist	Pedicle screw	Indirect	Cutting guide
Globus	Pedicle screw		Cutting guide

*TSolution One is manufactured by Think Surgical; Mako, by Stryker; NAVIO, by Smith & Nephew; ROSA, by Zimmer Biomet; OMNIBotics, by OMNI; SpineAssist, by MAZOR Robotics; and Globus, by Excelsius Medical. †THA = total hip arthroplasty, TKA = total knee arthroplasty, and UKA = unicondylar knee arthroplasty^{92,93}

not have a learning curve for accuracy in achieving the planned femoral and tibial implant position. Further, no additional risk for postoperative complications was observed during the surgical team learning curve.⁵⁵

A study by Jinnah et al. quantified the learning curve of robotic-arm assisted PKA. A total of 892 patients received a PKA performed by 11 different surgeons using the Mako System. Each surgeon had performed at least 30 surgeries with this technology, and the surgical time of the final 20 surgeries of each surgeon was averaged to define a steady state surgical time. The study measured the number of surgeries required to obtain two consecutive and three total surgeries completed within the 95% confidence interval of the steady state surgical time of that particular surgeon. Results showed that the number of surgeries required to have three surgeries completed within the 95% CI of the steady state surgical time was 13 (range: five to 29), and the number required to have two consecutive surgeries within this same time frame was 16 (range: four to 42).⁹⁵

Mako Total Knee studies have also shown a learning curve associated with Mako Total Knee before a surgical team can become time neutral to their operative time when performing manual TKA. Sodhi et al. performed a study to assess this learning curve in which two surgeons performed a total of 240 robotic-arm assisted cases.⁹⁶ Each case was allocated to a group of 20 sequential cases and a learning curve was created based on mean operative times. These times were compared to mean operative times for 20 randomly selected manual cases performed by the same surgeon. For Surgeon 1, mean operative time between the first and last cohort was reduced from 81 minutes to 70 minutes ($p < 0.05$). For Surgeon 2, mean operative time between the first and last cohort was reduced from 117 minutes to 98 minutes ($p < 0.05$). For both surgeons, the final 20-case set was time neutral to their manual cohort. This data implies that within a few months, a surgeon may be able to perform robotic-arm assisted TKA without any added operative time compared to manual TKA.⁹⁶

To understand how patient outcomes are influenced during a surgeon's learning curve, Sastry et al. reported on a single-surgeon experience comparing that surgeon's first 40 RATKA cases to a matched consecutive MTKA cohort.⁹⁷ During the first 40 cases, the RATKA cohort had a slightly greater overall surgical time when compared to the MTKA group (82.5 min vs. 78.3 min, $p=0.002$), however this difference was no longer statistically significant when only the second set of 20 RATKA cases

was considered (81.1 min vs. 78.3 min, $p = 0.254$). During this 40-case cohort, the RAKTA cohort showed a reduced LOS (1.27 days vs. 1.92 days, $p > 0.001$) and an improved ROM at 90 days ($+3.8^\circ$ vs. -8.7° , $p < 0.05$). No significant difference was noted in postoperative KSS or lower extremity activity scale at 30-, 60-, and 90-day follow-up between groups. The authors concluded that the surgeon's learning curve for RATKA appeared to progress rapidly, with a comparable OR time to MTKA by the second 20 cases.

Redmond et al. researched the learning curve during the adoption of RATHA as measured by component position, operative time and complications.⁹⁸ The first 105 robotic-arm assisted THAs performed by a single surgeon were divided into three groups based on the order of surgery: 1) Group A consisted of the first 35 patients who underwent Mako Total Hip by the senior surgeon, 2) Group B consisted of patients 36–70; and 3) Group C consisted of patients 71–105.⁹⁸ The authors reported a decreased risk of acetabular component malposition with Mako experience ($P < 0.05$).⁹⁸ Operative time appeared to decrease with increasing surgical experience with the Mako System ($P < 0.05$). A learning curve of 35 cases was observed with a decreased incidence of acetabular component outliers.

Heng and colleagues conducted a retrospective review of a single surgeon's last 45 conventional THAs performed prior to the surgeon's first 45 robotic-arm assisted THA. Surgical time, LOS in hospital, LOS in rehabilitation, transfusion rates and any complications were compared. The authors found that the average surgical time was 96.7 minutes for the robotic group and 84.9 minutes for the conventional group, however each robotic operation was approximately one minute shorter than the previous robotic operation and the average time for the last 10 robotic cases decreased to 82.9 mins.⁸⁴ Compared to conventional THA, there was no increased risk of complications or transfusions, and the authors noted there may be less chance of intraoperative acetabular fractures due to the single-ream, minimal bone resection technique utilized in the robotic procedure. LOS in the robotic group that did not go to rehabilitation was shorter by approximately one day and although a statistical analysis for LOS in rehabilitation was not performed due to small numbers, there was a tendency for shorter LOS in the robotic group as well.⁸⁴

Overall, the data showed that baseline operative times can be achieved, while increasing accuracy to plan.

The adoption of Mako SmartRobotics™

At the close of 2019, there were over 850 Mako Systems being used to perform surgery worldwide. Since launch, over 300,000 joint replacements have been performed with Mako. Additionally, 800 surgeons were trained on Mako Technology in the U.S. in 2019 alone.⁹⁹ The increasing adoption of Mako SmartRobotics™ Technology is supported by clinical success reported in published literature.

Mako has demonstrated the potential to deliver value through enhanced outcomes such as:


- Reduced pain and use of pain medications in TKA^{2,39,42}
- Reduced complications such as dislocation in THA, MUA in TKA and revision in UKA^{81,46,3}
- Increased patient satisfaction in TKA, THA and UKA^{42,83,3}
- In TKA and THA, reduced utilization of health services such as skilled nursing, home health aide, readmissions and emergency room^{48,49,51,89}
- Reduced payer cost in TKA, UKA and THA^{48,49,51,64}

Additionally, patients have reported benefits of Mako robotic-arm assisted procedures such as:

- Treatment satisfaction and return to activities of daily living^{81,83} for Mako Total Hip
- Treatment satisfaction^{3,59}, return to activities of daily living⁷³ and a “forgotten” joint^{56,60} for Mako Partial Knee
- Positive early outcomes measured using PROMs^{2,37,40,41} for Mako Total Knee; longer-term follow-up is ongoing

In summary, Mako SmartRobotics™ enables surgeons to achieve their target preoperative plans with precision, which may help distinguish them within their medical communities. The enhanced clinical outcomes observed to date with Mako SmartRobotics™ have the potential to provide value to patients, providers and payers alike.

14
years 
robotic-arm assisted
surgery experience

145+ 
published, peer
reviewed studies

 **850+**
Systems

have been installed across
26 countries and every state
in the contiguous U.S.*

 **1,000+**
U.S. and foreign patents
and patent applications
have been established

 **300K+**
Mako procedures
have been performed*



*Stryker's 2019 Q3 sales data

Figure 20. Mako SmartRobotics™ statistics⁹⁹

*Access to data analytics offering requires execution of a separate agreement. Availability of the data analytics offering may vary by country. Please speak with your sales representative for more information.

Cost-savings observed in the studies described in this document may differ across regions or countries due to differences in healthcare systems, treatment plans and associated costs.

References:

- Illgen RL, Bukowski BR, Abiola R, et al. Robotic-assisted total hip arthroplasty: outcomes at minimum two year follow up. *Surg Technol Int.* 2017;30:365-372.
- Kayani B, Konan S, Tahmassebi J, Pietrzak JRT, Haddad FS. Robotic-arm assisted total knee arthroplasty is associated with improved early functional recovery and reduced time to hospital discharge compared with conventional jig-based total knee arthroplasty: a prospective cohort study. *Bone Joint J.* 2018;100-B(7):930-937. doi:10.1302/0301-620X.100B7.BJJ-2017-1449.R1
- Kleeblad LJ, Borus T, Coon TM, Douchis J, Nguyen JT, Pearle AD. Midterm survivorship and patient satisfaction of robotic-arm assisted medial unicompartmental knee arthroplasty: a multicenter study. *J Arthroplasty.* 2018;33(6):1719-1726. doi:10.1016/j.arth.2018.01.0364
- Bell SW, Anthony I, Jones B, MacLean A, Rowe P, Blyth M. Improved accuracy of component positioning with robotic-assisted unicompartmental knee arthroplasty: data from a prospective, randomized controlled study. *J Bone Joint Surg Am.* 2016;98(8): 627-635. doi:10.2106/JBJS.15.00664
- Mahoney O, Kinsey T, Mont M, Hozack W, Orozco F, Chen A. Can computer generated 3D bone models improve the accuracy of total knee component placement compared to manual instrumentation? A prospective multi-center evaluation. Poster presented at: 32nd Annual Congress of the International Society for Technology in Arthroplasty (ISTA); October 2-5, 2019; Toronto, Canada.
- Banks SA. Haptic robotics enable a systems approach to design of a minimally invasive modular knee arthroplasty. *Am J Orthop (Belle Mead NJ).* 2009;38(2 Suppl):23-27.
- Suarez-Ahedo C, Gui C, Martin TJ, Chandrasekaran S, Lodhia P, Domb BG. Robotic-arm assisted total hip arthroplasty results in smaller acetabular cup size in relation to the femoral head size: a matched-pair controlled study. *Hip Int.* 2017;27(2):147-152. doi:10.5301/hipint.5000418
- Kayani B, Konan S, Pietrzak JRT, Haddad FS. Iatrogenic bone and soft tissue trauma in robotic-arm assisted total knee arthroplasty compared with conventional jig-based total knee arthroplasty: a prospective cohort study and validation of a new classification system. *J Arthroplasty.* 2018;33(8):2496-2501. doi:10.1016/j.arth.2018.03.042
- Hozack WJ. Multicentre analysis of outcomes after robotic-arm assisted total knee arthroplasty. *Bone Joint J:Orthop Proc.* 2018;100-B(Suppl_12):38.
- Hampp E, Chang TC, Pearle A. Robotic partial knee arthroplasty demonstrated greater bone preservation compared to robotic total knee arthroplasty. Poster presented at: Orthopaedic Research Society (ORS) Annual Meeting; February 2-5, 2019; Austin, TX.
- Sloan M, Premkumar A, Sheth N. Projected volume of primary total joint arthroplasty in the U.S., 2014 to 2030. *J Bone Joint Surg Am.* 2018;100(17):1455-1460. doi:10.2106/JBJS.17.01617
- Data and statistics. Centers for Disease Control and Prevention (CDC). Accessed April 22, 2020. https://www.cdc.gov/arthritis/data_statistics/index.htm
- Watkins-Castillo SI. Arthritis. The Burden of Musculoskeletal Diseases in the United States. Accessed April 22, 2020. <https://www.boneandjointburden.org/fourth-edition/iii0/arthritis>
- Barbour KE, Helmick CG, Theis KA, Murphy LB, Hootman JM, Brady TJ, Cheng YJ. Prevalence of doctor-diagnosed arthritis and arthritis-attributable activity limitation-United States, 2010-2012. *Morb Mortal Wkly Rep.* 2013; 62(44): 869-873.
- Cross M, Smith E, Hoy D, et al. The global burden of hip and knee osteoarthritis: estimates from the Global Burden of Disease 2010 Study. *Ann Rheum Dis.* 2014;73(7):1323-1330. doi:10.1136/annrheumdis-2013-204763
- Osteoarthritis: care and management clinical guideline [CG177]. National Institute for Health and Care Excellence (NICE). February 12, 2014. Accessed June 24, 2015. <https://www.nice.org.uk/guidance/cg177>
- Leopold SS. Minimally invasive total knee arthroplasty for osteoarthritis. *N Engl J Med.* 2009;360(17):1749-1758. doi:10.1056/NEJMct0806027
- Aleto TJ, Berend ME, Ritter MA, Faris PM, Meneghini RM. Early failure of unicompartmental knee arthroplasty leading to revision. *J Arthroplasty.* 2008;23(2):159-163. doi:10.1016/j.arth.2007.03.020
- National Joint Replacement Registry. Hip, Knee & Shoulder Arthroplasty Annual Report 2019. Australian Orthopaedic Association; 2019.
- Whiteside LA. Making your next unicompartmental knee arthroplasty last: three keys to success. *J Arthroplasty.* 2005;20(4 Suppl 2):2-3. doi:10.1016/j.arth.2005.03.029
- Ashraf T, Newman JH, Desai VV, Beard D, Nevelos JE. Polyethylene wear in a non-congruous unicompartmental knee replacement: a retrieval analysis. *Knee* 2004;11(3):177-181. doi:10.1016/j.knee.2004.03.004
- Hernigou P, Deschamps G. Posterior slope of the tibial implant and the outcome of unicompartmental knee arthroplasty. *J Bone Joint Surg Am.* 2004;86(3):506-511. doi:10.2106/00004623-200403000-00007
- Emerson RH Jr, Higgins LL. Unicompartmental knee arthroplasty with the oxford prosthesis in patients with medial compartment arthritis. *J Bone Joint Surg Am.* 2008;90(1):118-122. doi:10.2106/JBJS.F00739
- Suarez-Almazor ME, Richardson M, Kroll TL, Sharf BF. A qualitative analysis of decision-making for total knee replacement in patients with osteoarthritis. *J Clin Rheumatol.* 2010;16(4):158-163. doi:10.1097/RHU.0b013e3181df4de4
- Callaghan JJ, O'Rourke MR, Saleh KJ. Why knees fail: lessons learned. *J Arthroplasty.* 2004;19(4 Suppl 1):31-34. doi:10.1016/j.arth.2004.02.015
- Tarwala R, Dorr LD. Robotic assisted total hip arthroplasty using the MAKO platform. *Curr Rev Musculoskelet Med.* 2011;4(3):151-156. doi:10.1007/s12178-011-9086-7
- Hamilton DE, Burnett R, Patton JT, et al. Implant design influences patient outcome after total knee arthroplasty: a prospective double-blind randomised controlled trial. *Bone Joint J.* 2015;97-B(1):64-70. doi:10.1302/0301-620X.97B1.34254
- Mistry JB, Elmallah RK, Chughtai M, Oktem M, Harwin SE, Mont MA. Long-term survivorship and clinical outcomes of a single radius total knee arthroplasty. *Surg Technol Int.* 2016;28:247-251.
- Scott CEH, Clement ND, MacDonald DJ, et al. Five-year survivorship and patient-reported outcome of the Triathlon single-radius total knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc.* 2015;23(6):1676-1683. doi:10.1007/s00167-014-2922-8
- Wylde V, Blom AW, Whitehouse SL, et al. Patient-reported outcomes after total hip and knee arthroplasty: comparison of midterm results. *J Arthroplasty.* 2009;24(2):210-216. doi:10.1016/j.arth.2007.12.001
- Bourne RB, Chesworth BM, Davis AM, Mahomed NN, Charron KDJ. Patient satisfaction after total knee arthroplasty: who is satisfied and who is not? *Clin Orthop Relat Res.* 2010;468(1):57-63. doi:10.1007/s11999-009-1119-9
- Noble PC, Condit MA, Cook KF, Mathis KB. The John Insall Award: Patient expectations affect satisfaction with total knee arthroplasty. *Clin Orthop Relat Res.* 2006;452:35-43. doi:10.1097/01.blo.0000238825.63648.1e
- McNabb DC, Kim RH, Springer BD. Instability after total knee arthroplasty. *J Knee Surg.* 2015;28(2):97-104. doi:10.1055/s-0034-1396080
- Kim YH, Park JW, Kim J-S, Park S-D. The relationship between the survival of total knee arthroplasty and postoperative coronal, sagittal and rotational alignment of knee prosthesis. *Int Orthop.* 2014;38(2):379-385. doi:10.1007/s00264-013-2097-9
- Mason JB, Fehring TK, Estok R, Banel D, Fahrback K. Meta-analysis of alignment outcomes in computer-assisted total knee arthroplasty surgery. *J Arthroplasty.* 2007;22(8):1097-1106. doi:10.1016/j.arth.2007.08.001
- Khlopas A, Sodhi N, Sultan AA, Chughtai M, Molloy RM, Mont MA. Robotic arm-assisted total knee arthroplasty. *J Arthroplasty.* 2018;33(7):2002-2006. doi:10.1016/j.arth.2018.01.060
- Nickel B, Carroll K, Pearle A, Kleeblad LJ, Mayman DJ, Jerabek SA. Aim small, miss small: radiographic and functional outcomes of robotic-assisted total knee arthroplasty at one year. Presented at: 31st Annual Congress of the International Society for Technology in Arthroplasty (ISTA); October 10-13, 2018; London, United Kingdom.
- Bhowmik-Stoker M, Faizan A, Nevelos J, Tippett B, Clark G. Do total knee arthroplasty surgical instruments influence clinical outcomes? A prospective parallel study of 150 patients. Presented at: Orthopaedic Research Society (ORS) Annual Meeting; February 2-5, 2019; Austin, TX.
- Bhimani SJ, Bhimani R, Smith A, Eccles C, Smith L, Malkani A. Robotic-assisted total knee arthroplasty demonstrates decreased postoperative pain and opioid usage compared to conventional total knee arthroplasty. *Bone Joint Open.* 2020;1(2). doi:10.1302/2046-3758.12.BJO-2019-0004.R1
- Clark G. Australian Experience. Mako Robotic TKA. Presented at: Australian Orthopaedic Association (AOA) 77th Annual Scientific Meeting; October 8-12, 2017; Adelaide, Australia.
- Marchand RC, Sodhi N, Khlopas A, et al. Patient satisfaction outcomes after robotic arm-assisted total knee arthroplasty: a short-term evaluation. *J Knee Surg.* 2017;30(9):849-853. doi:10.1055/s-0037-1607450

42. Marchand RC, Sodhi N, Anis HK, et al. One-year patient outcomes for robotic-arm assisted versus manual total knee arthroplasty. *J Knee Surg.* 2019;32(11):1063-1068. doi:10.1055/s-0039-1683977
43. Marchand R, Marchand K, Taylor KB, Condrey CJ, Scholl L, Bhowmik-Stoker M. Patient satisfaction after robotic assisted total knee arthroplasty. Presented at: Annual International Society of Technology in Arthroplasty (ISTA) Meeting; October 2-5, 2019; Toronto, Canada.
44. Wang J, Mitchell J, Greiner J, Illgen R. Relative clinical outcomes comparing manual and robotic assisted total knee arthroplasty at minimum 1 year follow-up. Presented at: Orthopaedic Research Society (ORS) Annual Meeting, February 2-5, 2019. Austin, TX.
45. Malkani AL, Roche MW, Kolisek FR, et al. New technology for total knee arthroplasty provides excellent patient-reported outcomes: a minimum two-year analysis. *Surg Technol Int.* 2019;36:276-280.
46. Malkani AL, Roche MW, Kolisek FR, et al. Manipulation under anesthesia rates in technology-assisted versus conventional-instrumentation total knee arthroplasty. *Surg Technol Int.* 2020;36:336-340.
47. Hampf E, Scholl L, Faizan A, Westrich G, Mont M. Greater iatrogenic soft tissue damage in conventional approach when compared with the robotic-arm assisted approach for total knee arthroplasty. Poster presented at: 19th EFORT Congress; May 30-June 1, 2018; Barcelona, Spain.
48. Cool CL, Jacofsky DJ, Seeger KA, Sodhi N, Mont MA. A 90-day episode-of-care cost analysis of robotic-arm assisted total knee arthroplasty. *J Comp Eff Res.* 2019;8(5):327-336. doi:10.2217/ce-2018-0136
49. Mont MA, Cool C, Gregory D, Coppolecchia A, Sodhi N, Jacofsky DJ. Health care utilization and payer cost analysis of robotic arm assisted total knee arthroplasty at 30, 60, and 90 days. *J Knee Surg.* Accepted manuscript. Published online September 2, 2019. doi:10.1055/s-0039-1695741
50. McLawhorn AS, Buller LT. Bundled payments in total joint replacement: keeping our care affordable and high in quality. *Curr Rev Musculoskelet Med.* 2017;10(3):370-377. doi:10.1007/s12178-017-9423-6
51. Pierce J, Needham C, Adams C, Coppolecchia AB, Lavernia C. Robotic-assisted knee surgery: an economic analysis. Presented at: Orthopaedic Research Society Annual Meeting, February 8-11, 2020; Phoenix, AZ.
52. White RE Jr, Allman JK, Trauger JA, Dales BH. Clinical comparison of the midvastus and medial parapatellar surgical approaches. *Clin Orthop Relat Res.* 1999;OCt(367):117-122.
53. Tria AJ Jr, Coon TM. Minimal incision total knee arthroplasty: early experience. *Clin Orthop Relat Res.* 2003;Nov(416):185-190. doi:10.1097/01.blo.0000093030.56370.d9
54. Collier MB, Eickmann TH, Sukezaki F, McAuley JP, Engh GA. Patient, implant, and alignment factors associated with revision of medial compartment unicompartmental knee arthroplasty. *J Arthroplasty.* 2006;21(6 Suppl 2):108-115. doi:10.1016/j.arth.2006.04.012
55. Kayani D, Konan S, Pietrzak JRT, Huq SS, Tahmassebi J, Haddad FS. The learning curve associated with robotic arm assisted unicompartmental knee arthroplasty. *Bone Joint J.* 2018;100-B(8):1033-1042. doi:10.1302/0301-620X.100B8.BJJ-2018-0040.R1
56. Blyth MJG, Anthony I, Rowe P, Banger MS, MacLean A, Jones B. Robotic-arm assisted versus conventional unicompartmental knee arthroplasty: exploratory secondary analysis of a randomised controlled trial. *Bone Joint Res.* 2017;6(11):631-639. doi:10.1302/2046-3758.6.11.BJR-2017-0060.R1
57. Gilmour A, MacLean AD, Rowe PJ, Banger MS, Donnelly I, Jones BG, Blyth MJG. Robotic-arm-assisted vs conventional unicompartmental knee arthroplasty. The 2-year clinical outcomes of a randomized controlled trial. *J Arthroplasty.* 2018;33(7S):S109-S115. doi:10.1016/j.arth.2018.02.050
58. Kayani B, Konan S, Tahmassebi J, Rowan FE, Haddad FS. An assessment of early functional rehabilitation and hospital discharge in conventional versus robotic-arm assisted unicompartmental knee arthroplasty: a prospective cohort study. *Bone Joint J.* 2019;101-B(1):24-33. doi:10.1302/0301-620X.101B1.BJJ-2018-0564.R2
59. Pearle AD, van der List JP, Lee L, Coon TM, Borus TA, Roche MW. Survivorship and patient satisfaction of robotic-assisted medial unicompartmental knee arthroplasty at a minimum two-year follow-up. *Knee.* 2017;24(2):419-428. doi:10.1016/j.knee.2016.12.001
60. Zuiderbaan HA, Van der list JP, Khamaisy S, et al. Unicompartmental knee arthroplasty versus total knee arthroplasty: which type of artificial joint do patients forget? *Knee Surg Sports Traumatol Arthrosc.* 2015;25(3):681-686. doi:10.1007/s00167-015-3868-1
61. National Joint Replacement Registry. Hip, Knee & Shoulder Arthroplasty Annual Report 2017. Australian Orthopaedic Association; 2017.
62. Catani F. Clinical outcomes of robotically assisted UKAs at 3 years follow-up. Presented at: The Partial Knee Meeting; January 30-31, 2018; Bruges, Belgium.
63. National Joint Replacement Registry. Hip, Knee & Shoulder Arthroplasty Annual Report 2018. Australian Orthopaedic Association; 2018.
64. Cool CL, Needham KA, Khlopas A, Mont MA. Revision analysis of robotic arm-assisted and manual unicompartmental knee arthroplasty. *J Arthroplasty.* 2019;34(5):926-931. doi:10.1016/j.arth.2019.01.018
65. Deese JM, Gratto-Cox G, Carter DA, Sasser TM Jr, Brown KL. Patient reported and clinical outcomes of robotic-arm assisted unicompartmental knee arthroplasty: minimum two year follow-up. *J Orthop.* 2018;15(3):847-853. doi:10.1016/j.jor.2018.08.018
66. Dretakis K, Igoumenou VG. Outcomes of robotic-arm assisted medial unicompartmental knee arthroplasty: minimum 3-year follow-up. *Eur J Orthop Surg Traumatol.* 2019;29(6):1305-1311. doi:10.1007/s00590-019-02424-4
67. Banger M. 5 year survivorship of a randomized trial of robotic arm assisted vs manual uni-compartmental knee arthroplasty. Presented at: European Knee Society (EKS) Arthroplasty Conference; May 2-3, 2019; Valencia, Spain.
68. St Mart J-P, de Steiger RN, Cuthbert A, Donnelly W. The three-year survivorship of robotically assisted versus non-robotically assisted unicompartmental knee arthroplasty: a study from the Australian Orthopaedic Association National Joint Replacement Registry. *Bone Joint J.* 2020;102-B(3):319-328. doi:10.1302/0301-620X.102B3.BJJ-2019-0713.R1
69. Clement ND, Bell A, Simpson P, Macpherson G, Patton JT, Hamilton DF. Robotic-assisted unicompartmental knee arthroplasty has a greater early functional outcome when compared to manual total knee arthroplasty for isolated medial compartment arthritis. *Bone Joint Res.* 2020;9(1): 15-22. doi:10.1302/2046-3758.9.1.BJR-2019-0147.R1
70. Kazarian GS, Lonner JH, Maltenfort MG, Ghomrawi HMK, Chen AF. Cost-effectiveness of surgical and nonsurgical treatments for unicompartmental knee arthritis: a Markov Model. *J Bone Joint Surg Am.* 2018;100(19):1653-1660. doi:10.2106/JBJS.17.00837
71. Clement ND, Deehan DJ, Patton JT. Robot-assisted unicompartmental knee arthroplasty for patients with isolated medial compartment osteoarthritis is cost-effective: a Markov decision analysis. *Bone Joint J.* 2019;101-B(9):1063-1070. doi:10.1302/0301-620X.101B9.BJJ-2018-1658.R1
72. Crawford R, Varughese I, Herron E, Jaiprakash A, Donnelly W, Whitehouse S. The cost effectiveness of unicompartmental knee arthroplasty vs total knee arthroplasty. Presented at: 79th Annual Scientific Meeting of the Australian Orthopaedic Association (AOA); October 6-10, 2019; Canberra, Australia.
73. Borus T, Roberts D, Fairchild P, Pirtle K, Baer M. Early functional performance of unicompartmental knee arthroplasty compared to total knee arthroplasty. Presented at: 2nd World Arthroplasty Congress (WAC); April 19-21, 2018. Rome, Italy.
74. Brown NM, Sheth NE, Davis K, et al. Total knee arthroplasty has higher postoperative morbidity than unicompartmental knee arthroplasty: a multicenter analysis. *J Arthroplasty.* 2012;27(8 Suppl):86-90. doi:10.1016/j.arth.2012.03.022
75. Knight SR, Aujla R, Biswas SP. Total hip arthroplasty - over 100 years of operative history. *Orthop Rev (Pavia).* 2011;3(2):e16. doi:10.4081/or.2011.e16
76. Callanan MC, Jarrett B, Bragdon CR, et al. The John Charnley Award: risk factors for cup malpositioning: quality improvement through a joint registry at a tertiary hospital. *Clin Orthop.* 2011;469(2):319-329. https://doi.org/10.1007/s11999-010-1487-1
77. Bozic KJ, Kurtz SM, Lau E, Ong K, Vail TP, Berry DJ. The epidemiology of revision total hip arthroplasty in the United States. *J Bone Joint Surg Am.* 91(1):128-133. doi:10.2106/JBJS.H.00155
78. Elson L, Dounchis J, Illgen R, et al. Precision of acetabular cup placement in robotic integrated total hip arthroplasty. *Hip Int.* 2015;25(6):531-536. doi:10.5301/hipint.5000289
79. Domb BG, Redmond JM, Louis SS, et al. Accuracy of component positioning in 1980 total hip arthroplasties: a comparative analysis by surgical technique and mode of guidance. *J Arthroplasty.* 2015;30(12):2208-2218. doi:10.1016/j.arth.2015.06.059
80. Domb BG, El Bitar YF, Sadik AY, Stake CE, Botser IB. Comparison of robotic-assisted and conventional acetabular cup placement in THA: a matched-pair controlled study. *Clin Orthop Relat Res.* 2014;472(1):329-336. doi:10.1007/s11999-013-3253-7
81. Bukowski BR, Anderson P, Khlopas A, Chughtai M, Mont MA, Illgen RL. Improved functional outcomes with robotic compared with manual total hip arthroplasty. *Surg Technol Int.* 2016;29:303-308.
82. Perets I, Walsh JP, Close MR, Mu BH, Yuen LC, Domb BG. Robot-assisted total hip arthroplasty: clinical outcomes and complication rate. *Int J Med Robot; 2018;14(4):e1912.* doi:10.1002/rcs.1912

83. Domb BG, Chen JW, Lall AC, Perets I, Maldonado DR. Minimum 5-year outcomes of robotic-assisted primary total hip arthroplasty with a nested comparison against manual primary total hip arthroplasty: a propensity score-matched study. *J Am Acad Orthop Surg*. Accepted manuscript. Published online February 25, 2020. doi:10.5435/JAAOS-D-19-00328
84. Heng YY, Gunaratne R, Ironside C, Taheri A. Conventional vs robotic-arm assisted total hip arthroplasty (THA) surgical time, transfusion rates, length of stay, complications and learning curve. *J Arthritis*. 2018;7(4):1-4. doi:10.4172/2167-7921.1000272
85. Mehra A, Moriso ZN, Schemitsch E, Waddell J. Assessing leg lengths intraoperatively in total hip arthroplasty: comparison of two methods. *Surg Technol Int*. 2013;23:258-260.
86. Manzotti A, Cerveri P, De Momi E, Pullen C, Confalonieri N. Does computer-assisted surgery benefit leg length restoration in total hip replacement? navigation versus conventional freehand. *Int Orthop*. 2011;35(1):19-24. doi:10.1007/s00264-009-0903-1
87. Xu B, Yang D, Aili R, Cao L. Effect of femoral offset change on pain and function after total hip arthroplasty. *Zhongguo Xiu Fu Chong Jian Wai Ke Za Zhi*. 2013;27(7):843-846.
88. Nodzo SR, Chang C-C, Carroll KM, et al. Intraoperative placement of total hip arthroplasty components with robotic-arm assisted technology correlates with postoperative implant position: a CT-based study. *Bone Joint J*. 2018;100-B(10):1303-1309. doi:10.1302/0301-620X.100B10-BJJ-2018-0201.R1
89. Pierce J, Needham K, Adams C, Coppolecchia A, Lavernia C. Robotic assisted total hip arthroplasty: a 90-Day episode of care cost analysis. Presented at: The Professional Society for Health Economics and Outcomes Research (ISPOR) Annual Meeting; May 18-20, 2020; Virtual.
90. Jacofsky DJ, Allen M. Robotics in arthroplasty: a comprehensive review. *J Arthroplasty*. 2016;31(10):2353-2363. doi:10.1016/j.arth.2016.05.026
91. Mezger U, Jendrewski C, Bartels M. Navigation in surgery. *Langenbecks Arch Surg*. 2013;398(4):501-514. doi:10.1007/s00423-013-1059-4
92. Chen AF, Kazarian GS, Jessop GW, Makhdom A. Robotic technology in orthopaedic surgery. *J Bone Joint Surg Am*. 2018;100(22):1984-1992. doi:10.2106/JBJS.17.01397
93. Mathew KK, Marchard KB, Tarazi JM, et al. Computer-assisted navigation in total knee arthroplasty. *Surg Technol Int*. 2020;36:323-330.
94. Sultan AA, Samuel LT, Khlopas A, et al. Robotic-arm assisted total knee arthroplasty more accurately restored the posterior condylar offset ratio and the Insall-Salvati Index compared to the manual technique; a cohort-matched study. *Surg Technol Int*. 2019;34:409-413.
95. Jinnah R, Lippincott CJ, Horowitz S, Conditt MA. The learning curve of robotically-assisted UKA. Presented at: Orthopaedic Research Society (ORS) 56th Annual Meeting; March 6-10, 2010; New Orleans, LA.
96. Sodhi N, Khlopas A, Piuizzi NS, et al. The learning curve associated with robotic total knee arthroplasty. *J Knee Surg*. 2018;31(1):17-21. doi:10.1055/s-0037-1608809
97. Naziri Q, Cusson BC, Chaudhri M, Shah NV, Sastry A. Making the transition from traditional to robotic-arm assisted TKA: what to expect? A single-surgeon comparative-analysis of the first 40-consecutive cases. *J Orthop*. 2019;16(4):364-368. doi: 10.1016/j.jor.2019.03.010
98. Redmond JM, Gupta A, Hammarstedt JE, Petrakos AE, Finch NA, Domb BG. The learning curve associated with robotic-assisted total hip arthroplasty. *J Arthroplasty*. 2015;30(1):50-54. doi:10.1016/j.arth.2014.08.003
99. Data on file. Stryker sales data, 2019.
100. Conditt M, Coon T, Roche M, et al. Short to mid term survivorship of robotically assisted UKA: a multicenter study. Presented at: International Society for Technology in Arthroplasty (ISTA) Annual Meeting; September 24-27, 2014; Kyoto, Japan.
101. Burger JA, Kleeblad LJ, Laas N, Pearle AD. Mid-term survivorship and patient-reported outcomes of robotic-arm assisted partial knee arthroplasty. *Bone Joint J*. 2020;102-B(1):108-116. doi:10.1302/0301-620X.102B1.BJJ-2019-0510.R1

A surgeon must always rely on his or her own professional clinical judgment when deciding whether to use a particular product when treating a particular patient. Stryker does not dispense medical advice and recommends that surgeons be trained in the use of any particular product before using it in surgery.

The information presented is intended to demonstrate the breadth of Stryker's product offerings. A surgeon must always refer to the package insert, product label and/or instructions for use before using any of Stryker's products. The products depicted are CE marked according to the Medical Device Regulation 2017/745 or the Medical Device Directive 93/42/EEC. Products may not be available in all markets because product availability is subject to the regulatory and/or medical practices in individual markets. Please contact your sales representative if you have questions about the availability of products in your area.

Stryker Corporation or its divisions or other corporate affiliated entities own, use or have applied for the following trademarks or service marks: AccuStop, Mako, SmartRobotics, Stryker. All other trademarks are trademarks of their respective owners or holders.

MKOEVS-AR-2_25812

Copyright © 2020 Stryker

Howmedica Osteonics Corp.

325 Corporate Drive

Mahwah, NJ 07430, USA

A subsidiary of Stryker Corporation

t: 201 831 5000

stryker.com